



## Data in Brief

# Transcriptome analysis of T<sub>H</sub>2 CD4<sup>+</sup> T cells differentiated from wild-type and NLRP3KO mice



Mélanie Bruchard <sup>a</sup>, Romain Boidot <sup>a,b,\*</sup>, François Ghiringhelli <sup>a,b</sup>, Frédérique Végran <sup>a,b</sup>

<sup>a</sup> Inserm U866, Dijon, France

<sup>b</sup> Centre Georges-François Leclerc, Platform of Transfer in Cancer Biology, Dijon, France

## ARTICLE INFO

## Article history:

Received 18 June 2015

Received in revised form 24 June 2015

Accepted 24 June 2015

Available online 9 July 2015

## Keywords:

Transcriptome

Th2 differentiation

NLRP3

## ABSTRACT

The Nod-like receptor NLRP3 is involved in the formation of inflammasome. Up to now, the immunological functions of NLRP3 independently of inflammasome is unclear. In this dataset containing 6 samples (T<sub>H</sub>0, T<sub>H</sub>2 cells at day 3 and day 6 in wild type or *Nlrp3* deficient cells), we show that NLRP3 expression in CD4<sup>+</sup> T cells supports a T helper 2 (T<sub>H</sub>2) transcriptional program in a cell-intrinsic manner (raw and normalized data are accessible on Gene Expression Omnibus database under the number GSE54561, <http://www.dtd.nlm.nih.gov/geo/query/acc.cgi?acc=GSE54561>). Indeed, NLRP3 positively-regulated T<sub>H</sub>2 program independently of inflammasome formation. These data indicated that T<sub>H</sub>2 specific genes such as cMaf or IL4 were not induced in *Nlrp3* deficient cells. These results demonstrate the capacity of NLRP3 to act as a key transcription factor in T<sub>H</sub>2 differentiation.

© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Specifications	
Organism/cell line/tissue	<i>Mus musculus</i>
Sex	Sex independent
Sequencer or array type	MiSeq (Illumina)
Data format	Raw data and normalized data
Experimental factors	T CD4 lymphocytes obtained from mice and differentiated in Th2 subtype
Experimental features	The transcriptional profile of T CD4 lymphocytes obtained from wild-type or NLRP3KO mice differentiated in Th2 cells was analyzed by RNA sequencing.
Consent	Protocol approved by ethic committee
Sample source location	France

## 1. Direct link to deposited data

<http://www.dtd.nlm.nih.gov/geo/query/acc.cgi?acc=GSE54561>.

## 2. Experimental design, materials and methods

The transcriptome of CD4<sup>+</sup> T cells was studied by RNA sequencing on MiSeq device (Illumina). Naive CD4<sup>+</sup> T cells (CD4<sup>+</sup>CD62L<sup>high</sup>) from spleens and lymph nodes of C57BL/6 *Nlrp3* wild type or *Nlrp3*<sup>−/−</sup> mice

were sorted by flow cytometry with BD ARIA cytometer equipped with BD FACSDiva software (BD Biosciences). The purity of isolated naive T cell population exceeded 95%. Naive T cells were cultured for three days with anti-CD3 (2 µg/mL) and anti-CD28 (2 µg/mL) antibodies in the presence of anti-IFNγ (10 µg/mL) and anti-IL-4 (10 µg/mL) to obtain T<sub>H</sub>0 or of anti-IL-4 (10 µg/mL) and IL-12 (10 ng/mL) to obtain T<sub>H</sub>1 or of anti-IFNγ (10 µg/mL) and IL-4 (10 ng/mL) to obtain T<sub>H</sub>2. Cells were cultured at 37 °C under 5% CO<sub>2</sub> in RPMI 1640 with 5% (vol/vol) fetal calf serum supplemented with sodium pyruvate, penicillin and streptomycin, and 4 mM of 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES). Anti-CD3 (clone 145-2C11), anti-CD28 (clone PV-1), anti-IL-4 (clone 1B11) and anti-IFN-γ (clone XMG.1) antibodies were obtained from BioXcell and IL-12 and IL-4 were purchased from R&D Systems. A second round of stimulation was done using the same cytokine cocktails plus IL-2 (100 U/mL, R&D Systems) for another three days.

For RNA-Seq library preparation, total RNA from T cells was extracted using TRIzol (Life Technologies). rRNA was removed, thanks to Ribominus Kit (Life Technologies). 100 ng of rRNA depleted RNA was used for the library preparation, thanks to the NEBNext Ultra RNA library kit for Illumina (New England Biolabs) by following the manufacturer's instructions. Briefly, RNA was heat fragmented and a reverse transcription was performed. Next, the second strand of cDNA was synthesized and the reaction mixture was purified with magnetic beads. Afterwards, an end repair step was carried out and adapters were ligated. After second magnetic bead purification, a PCR of 12 cycles was performed to increase the target yield and incorporate index. Quality and quantity of libraries were assessed after magnetic bead purification thanks to a TapeStation fragment analyzer (Agilent). Library size was

\* Corresponding author at: Centre Georges-François Leclerc, Platform of Transfer in Cancer Biology, Dijon, France.

between 250 base pairs and 400 base pairs. Each library was diluted at a concentration of 4 nM. Five libraries were pooled and 12 pM of the mix was injected into the sequencing flow cell. The RNA-Seq libraries were sequenced with paired-end 151 bp reads. Data analysis was carried out via Galaxy interface [1]. FASTQ files were mapped by using BWA (mm9 version of *Mus musculus* genome) for Illumina [2]. The analysis was performed by using TopHat for Illumina [3]. Generated files were processed with Cufflinks software [4] to obtain annotated expressed genes in each studied subtype. Then, differential expression between the samples was analyzed with Cuffdiff [4]. Raw and normalized data were accessible on public database: GEO submission number GSE54561, <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE54561> [5].

## References

- [1] J. Taylor, I. Schenck, D. Blankenberg, A. Nekrutenko, Using galaxy to perform large-scale interactive data analyses. *Curr. Protoc. Bioinformatics* (2007) (Chapter 10: Unit 10 15).
- [2] H. Li, R. Durbin, Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* 25 (2009) 1754–1760.
- [3] C. Trapnell, L. Pachter, S.L. Salzberg, TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* 25 (2009) 1105–1111.
- [4] C. Trapnell, B.A. Williams, G. Pertea, A. Mortazavi, G. Kwan, M.J. van Baren, et al., Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* 28 (2010) 511–515.
- [5] M. Bruchard, C. Reb  , V. Derang  re, D. Togb  , B. Ryffel, R. Boidot, E. Humblin, E. Hamman, F. Chalmin, H. Berger, A. Chevriaux, E. Limagne, L. Apetoh, F. V  gran, F. Ghiringhelli, The Nod-like receptor NLRP3 is a transcriptional regulator of T helper 2 cell differentiation. *Nat. Immunol.* (2015) (in press).